

Claims:

1. Method for forming carbon-carbon bonds by reacting compounds of the general formula type (I)



- 5 with π compounds in neutral to basic aqueous or alcoholic solvents or solvent mixtures which have a lower nucleophilicity than the π nucleophile,

where R^1 , R^2 and R^3 are independently of one another organic or organometallic radicals or hydrogen,

- 10 where X is a leaving group of a type such that the compounds (I) are solvolysed in ethanol with a rate constant k_{EtOH} of $> 10^{-6} \text{ s}^{-1}$ (25°C),

characterized in that the intermediate carbocations are generated in neutral to basic aqueous or alcoholic solvents or solvent mixtures without using a Lewis acid or protic acid.

- 15 2. Method according to Claim 1, characterized in that R^1 , R^2 and R^3 are chosen independently of one another from the group of branched or unbranched alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, branched or unbranched alk-2-enyl, cyclo-, bicyclo- and tricycloalkyl, alkoxy, aryloxy or hydrogen, where two of the radicals R^1 , R^2 and R^3 form an alkyl ring where appropriate.

- 20 3. Method according to either of the preceding claims, characterized in that X is a leaving group chosen from the group of halogen, alkoxy, alkyl- or arylsulphonato, substituted or unsubstituted phenoxy, acyloxy, benzoyloxy, carbamoyl, alkyloxycarbonyloxy, aryloxycarbonyloxy, siloxy, phosphato, phosphonato, hypophosphonato, alkylperoxy, sulphato, sulphenyl, sulphonyl, S-alkylsulphoxy, S-arylsulphoxy, alkylthio, arylthio, thiocyanato, isothiocyano, ureato and imidyl.
- 25

4. Method according to any of the preceding claims, characterized in that the solvent or solvent mixture used comprises solvent components from the group of water, ethanol, methanol, 2,2,2-trifluoroethanol, 1,1,1,3,3,3-hexafluoro-2-propanol, tetrahydrofuran, acetone, acetonitrile and dioxane.
- 5 5. Method according to any of the preceding claims, characterized in that compounds of type (I) are reacted with a mixture of a π compound in the appropriate solvent or solvent mixture and possibly further, basic inorganic or organic additions.
- 10 6. Method according to any of the preceding claims, characterized in that the π compounds used as nucleophile are aliphatic π compounds from the group of substituted alkenes and alkynes, allyl- and propargylsilanes, alkyl enol ethers, silyl enol ethers, (silyl) ketene acetals and enamines, or aromatic π compounds from the group of donor-substituted or unsubstituted aromatic and heteroaromatic compounds.